arXiv:1207.5335v1 [math.PR] 23 Jul 2012

The Invariant Measure of Random Walks in the Quarter-plane: Representation in Geometric Terms

Yanting Chen*, Richard J. Boucherie*, and Jasper Goseling*

*Stochastic Operations Research, Department of Applied Mathematics, University of Twente

July 24, 2012

Abstract

We consider the invariant measure of homogeneous random walks in the quarter-plane. In particular, we consider measures that can be expressed as a finite or countable linear combination of geometric terms. We present conditions on the cardinality and the structure of these linear combinations such that the resulting measure can be an invariant measure of a random walk. Our contributions are as follows: First, we demonstrate that each geometric term must individually satisfy the balance equations in the interior of the state. Next, it is shown that the geometric terms in an invariant measure must have a pairwise-coupled structure. Finally, it is shown that a linear combination of two geometric terms can never be an invariant measure. Moreover, if a finite linear combination of geometric terms is an invariant measure, then at least one coefficient must be negative.

Keywords: random walk, quarter-plane, positive orthant, invariant measure, stationary distribution, product-form, compensation method, geometric distribution.

---

*P.O. Box 217, 7500 AE Enschede, The Netherlands. Email:{Y.Chen, R.J.Boucherie, J.Goseling}@utwente.nl

1 Introduction

We study random walks in the quarter-plane that are homogeneous in the sense that transition probabilities are translation invariant. Our interest is in invariant measures that can be expressed as linear combinations of geometric terms, *i.e.*, the measure m in state (i, j) is of the form

$$m(i,j) = \sum_{(\rho,\sigma)} \alpha(\rho,\sigma)\rho^i\sigma^j.$$

Many examples exist of random walks with geometric product-form invariant measure, *i.e.*, with a single geometric term, see *e.g.*, [3]. In addition, it is known that the invariant measure of some random walks can be expressed as a linear combination of countably many geometric terms [2]. This naturally leads to the problem addressed in the current work, that is to characterize which linear combinations of geometric measures may yield invariant measures for the random walk in the quarter-plane.

Various approaches to finding the invariant measure of a random walk in the quarter-plane exist. Most notably, methods from complex analysis have been used for finding expressions for the generating function of the invariant measure for arbitrary random walks [5, 7]. Matrix-geometric methods provide an algorithmic approach to finding the invariant measure [9]. However, explicit expressions for the invariant measures of most random walks are still hard to obtain using these methods. An overview of recent work on finding the invariant measure of random walks in the quarter-plane is given in [8].

The focus of the current work is on the class of measures that can be expressed as a finite or countable linear combination of geometric terms. We present conditions on the cardinality and the structure of these linear combinations such that the resulting measure can be an invariant measure of a random walk. Our contributions are as follows: First we demonstrate that each geometric term must individually satisfy the balance equations in the interior of the state. Next it is shown that the geometric terms in an invariant measure must have a pairwise-coupled structure. Finally, it is shown that a linear combination of two geometric terms can never be an invariant measure. Moreover, if a finite linear combination of geometric terms is an invariant measure, then at least one coefficient must be negative. A similar study for processes on a continuous state space, *i.e.*, for reflected Brownian motion, is presented in [6].

The remainder of this paper is structured as follows. In Section 2 we present the model. Definitions and some initial results are given in Section 3. Necessary conditions on the structure of the set of geometric terms are given in Section 4. In Section 5 conditions on the cardinality of this set and the signs of the coefficients in the linear combination are investigated.

2 Model

Consider a two-dimensional random walk on the pairs (i,j) of non-negative integers. We refer to $\{(i,j)|i>0, j>0\}$, $\{(i,j)|i>0, j=0\}$, $\{(i,j)|i=0, j>0\}$ and $(0,0)$ as the interior, the horizontal axis, the vertical axis and the origin of the state space, respectively. The transition probability from state (i,j) to state $(i+s, j+t)$ is denoted by $p_{s,t}(i,j)$. Transitions are restricted to the adjoining points (horizontally, vertically and diagonally), *i.e.*, $p_{s,t}(k,l)=0$ if $|s|>1$ or $|t|>1$. The process is homogeneous in the sense that for each pair (i,j), (k,l) in the interior (respectively on the horizontal axis and on the vertical axis) of the state space

$$p_{s,t}(i,j) = p_{s,t}(k,l) \quad \text{and} \quad p_{s,t}(i-s,j-t) = p_{s,t}(k-s,l-t), \tag{1}$$

for all $-1 \leq s \leq 1$ and $-1 \leq t \leq 1$. We introduce, for $i>0$, $j>0$, the notation $p_{s,t}(i,j)=p_{s,t}$, $p_{s,0}(i,0)=h_s$ and $p_{0,t}(0,j)=v_t$. Note that the first equality of (1) implies that the transition probabilities for each part of the state space are translation invariant. The second equality ensures that also the transition probabilities entering the same part of the state space are translation invariant.[1] The above definitions imply that $p_{1,0}(0,0)=h_1$ and $p_{0,1}(0,0)=v_1$. The model and notation are illustrated in Figure 1.

We assume that the random walk is ergodic and has invariant measure

---

[1] Note that this is a stronger notion of homogeneity than considered in, for instance, [2].



Figure 1: Random walk in the quarter-plane.

m, *i.e.*, for $i > 0$ and $j > 0$

$$m(i,j) = \sum_{s=-1}^{1} \sum_{t=-1}^{1} m(i-s, j-t) p_{s,t}, \tag{2}$$

$$m(i,0) = \sum_{s=-1}^{1} m(i-s,1) p_{s,-1} + \sum_{s=-1}^{1} m(i-s,0) p_{s,0}, \tag{3}$$

$$m(0,j) = \sum_{t=-1}^{1} m(1,j-t) p_{-1,t} + \sum_{t=-1}^{1} m(0,j-t) p_{0,t}. \tag{4}$$

We will refer to the above as the balance equations in the interior, the horizontal axis and the vertical axis of the state space respectively.

3 Preliminaries

In the first part of this section we show that for invariant measures consisting of linear combinations of geometric terms, these geometric terms individually satisfy the interior balance equations. In the second part of the section we introduce the notions of uncoupled partitions and pairwise coupled sets which will be used in later sections.



(a) (b) (c) (d)

Figure 2: Examples of $Q(\rho,\sigma)=0$. In 2(a) $p_{1,0}=p_{0,1}=\frac{1}{5}$, $p_{-1,-1}=\frac{3}{5}$. In 2(b) $p_{1,0}=\frac{1}{5}=p_{0,-1}=\frac{1}{5}$, $p_{-1,1}=\frac{2}{5}$. In 2(c) $p_{1,1}=\frac{1}{62}, p_{-1,1}=p_{1,-1}=\frac{10}{31}, p_{-1,-1}=\frac{21}{62}$. In 2(d) $p_{-1,1}=p_{1,-1}=\frac{1}{4}, p_{-1,-1}=\frac{1}{2}$.

3.1 Induced measures

Definition 1 (Induced measure). *Measure m is called induced by $\Gamma \subset \mathbb{R}_+^2$ if*

$$m(i,j)=\sum_{(\rho,\sigma)\in\Gamma}\alpha(\rho,\sigma)\rho^i\sigma^j,$$

with $\alpha(\rho,\sigma)\neq 0$ for all $(\rho,\sigma)\in\Gamma$.

We restrict ourselves to finite measures, for which the value of the summation is independent of the ordering of the sequences. In other words, we assume that $\sum_{(\rho,\sigma)\in\Gamma}|\alpha(\rho,\sigma)|(1-\rho)^{-1}(1-\sigma)^{-1}<\infty$. Moreover, we restrict our attention to Γ of at most countably infinite cardinality. In particular, we will consider linear combinations of geometric measures that individually satisfy the balance equations in the interior of the state space, (2). We introduce the polynomial

$$Q(\rho,\sigma)=\rho\sigma\left(\sum_{s=-1}^{1}\sum_{t=-1}^{1}\rho^{-s}\sigma^{-t}p_{s,t}-1\right) \tag{5}$$

to capture the notion of balance, *i.e.*, $Q(\rho,\sigma)=0$ implies that $\rho^i\sigma^j$ satisfies (2). Several examples of $Q(\rho,\sigma)=0$ can be found in Figure 2. Let C be the restriction of $Q(\rho,\sigma)=0$ to the interior of the positive unit square, *i.e.*,

$$C=\Big\{(\rho,\sigma)\in(0,1)^2 \mid Q(\rho,\sigma)=0\Big\}. \tag{6}$$

The next result justifies the restriction of considering only pairs (ρ,σ) from C.

Theorem 1. *If the invariant measure for a random walk in the quarter-plane is induced by* $\Gamma \subset \mathbb{R}_+^2$ *where* Γ *is of at most countable cardinality, then* $\Gamma \subset C$.

Proof. Since m satisfies balance in the interior of the state space

$$\sum_{k=1}^{\infty} \rho_k^i \sigma_k^j [\alpha(\rho_k, \sigma_k)(1 - \sum_{s=-1}^{1} \sum_{t=-1}^{1} \rho_k^{-s} \sigma_k^{-t} p_{s,t})] = 0. \tag{7}$$

Moreover, we have

$$\begin{aligned} &\sum_{k=1}^{\infty} \alpha(\rho_k, \sigma_k)\rho_k \sigma_k (1 - \sum_{s=-1}^{1} \sum_{t=-1}^{1} \rho_k^{-s} \sigma_k^{-t} p_{s,t}) \\ <&M \sum_{k=1}^{\infty} |\alpha(\rho_k, \sigma_k)| \frac{1}{1-\rho} \frac{1}{1-\sigma} \\ <&\infty, \end{aligned}$$

where M is a finite positive constant. Hence, the sequence $\alpha(\rho_k, \sigma_k)\rho_k\sigma_k(1 - \sum_{s=-1}^{1} \sum_{t=-1}^{1} \rho_k^{-s} \sigma_k^{-t} p_{s,t})$, where $k = 1, 2, \cdots$, is absolutely convergent. Then, it follows directly from [4, Theorem 1] that

$$1 - \sum_{s=-1}^{1} \sum_{t=-1}^{1} \rho_k^{-s} \sigma_k^{-t} p_{s,t} = 0.$$

which is the interior balance equation for the measure $\rho_k^i \sigma_k^j$, hence $Q(\rho_k, \sigma_k) = 0$. Finally note that an element $(\rho, \sigma) \notin (0, 1)^2$ would induce a measure that is not finite. □

3.2 Uncoupled partitions and pairwise-coupled sets

Different ways of partitioning set Γ will be introduced here. These partitions play an essential role in the analysis later on.

Definition 2 (Uncoupled partition)**.** *A partition* $\{\Gamma_1, \Gamma_2, \cdots\}$ *of* Γ *is* horizontally uncoupled *if* $(\rho, \sigma) \in \Gamma_i$ *and* $(\tilde{\rho}, \tilde{\sigma}) \in \Gamma_j$ *for* $i \neq j$, *implies that* $\tilde{\rho} \neq \rho$; *is* vertically uncoupled *if* $(\rho, \sigma) \in \Gamma_i$ *and* $(\tilde{\rho}, \tilde{\sigma}) \in \Gamma_j$ *for* $i \neq j$, *implies that* $\tilde{\sigma} \neq \sigma$; *and is* uncoupled *if it is both horizontally and vertically uncoupled.*



(a) (b) (c) (d)

Figure 3: Partitions of Γ. In (a) an example of curve C and subset Γ are given. Figures (b), (c) and (d) give the uncoupled, horizontally uncoupled and vertically uncoupled partitions of Γ respectively, where different sets are marked by different symbols.

We call the partition with the largest number of sets a maximal partition.

Lemma 1. *The maximal horizontally uncoupled partition is unique.*

Proof. Assume that $\{\Gamma_i\}_{i=1}^H$ and $\{\Gamma'_i\}_{i=1}^H$ are different maximal horizontally uncoupled partitions of Γ. Without loss of generality, $\Gamma_1 \cap \Gamma'_1 \neq \emptyset$ and $\Gamma_1 \setminus \Gamma'_1 \neq \emptyset$. Consider $(\rho, \sigma) \in \Gamma_1 \setminus \Gamma'_1$ and $(\tilde{\rho}, \tilde{\sigma}) \in \Gamma_1 \cap \Gamma'_1$. If $\rho = \tilde{\rho}$, then $\{\Gamma'_i\}_i$ is not a horizontally uncoupled partition. If $\rho \neq \tilde{\rho}$, then $\{\Gamma_i\}_i$ is not maximal. $\square$

Existence of unique maximal (vertically) uncoupled partitions follows similarly. Examples of a maximal uncoupled partition, of a maximal horizontally uncoupled partition and of a maximal vertically uncoupled partition can be found in Figures 3(b), 3(c) and 3(d) respectively.

Let H denote the number of elements in the maximal horizontally uncoupled partition and Γ_i^h, $i = 1, \ldots, H$, the sets themselves. The common horizontal coordinate of set Γ_i^h is denoted by $\varrho(\Gamma_i^h)$. The maximal vertically uncoupled partition has V sets, Γ_j^v, $j = 1, \cdots, V$, where elements of Γ_j^v have common vertical coordinate $\varsigma(\Gamma_j^v)$.The maximal uncoupled partition is denoted by $\{\Gamma_k^u\}_{k=1}^U$.

Next, we make two observations on the structure of $\Gamma \subset C$ for which the maximal uncoupled partition consists of only one set. Firstly, for any $(\rho, \sigma) \in \Gamma$ there always exist either $(\rho, \tilde{\sigma}) \in \Gamma$, $\tilde{\sigma} \neq \sigma$ or $(\tilde{\rho}, \sigma) \in \Gamma$, $\tilde{\rho} \neq \rho$. Secondly, the degree of $Q(\rho, \sigma)$ is at most two in each variable. This means, for instance, that if $(\rho, \sigma) \in \Gamma$ and $(\rho, \tilde{\sigma}) \in \Gamma$, $\tilde{\sigma} \neq \sigma$, then there does not exist $(\rho, \hat{\sigma}) \in \Gamma$, where $\hat{\sigma} \neq \sigma$, $\hat{\sigma} \neq \tilde{\sigma}$ and $\hat{\sigma} \neq \sigma$. By repeating the above two



(a) (b) (c) (d)

Figure 4: Examples of pairwise-coupled sets.

arguments for other elements in Γ it follows that Γ must have a pairwise-coupled structure. An example of such a set is $\Gamma = \{(\rho_k, \sigma_k), k = 1, 2, 3 \cdots\}$ where

$$\rho_1 = \rho_2, \sigma_1 > \sigma_2, \rho_2 > \rho_3, \sigma_2 = \sigma_3, \rho_3 = \rho_4, \sigma_3 > \sigma_4, \cdots . \tag{8}$$

The above discussion leads to the definition of a pairwise-coupled set in terms of the number of sets in a maximal uncoupled partition.

Definition 3 (Pairwise-coupled)**.** *A set $\Gamma \subset C$ is called pairwise-coupled if and only if the maximal uncoupled partition of Γ contains only one set.*

Some examples of pairwise-coupled sets can be found in Figure 4. The example of countably many pairwise-coupled elements given in (8) occurs in, for instance, [2] when applying the compensation method. A sufficient condition for applying the compensation method is that the random walk does not have transition probabilities to the North, North-East and East, *i.e.*, that $p_{1,0} = p_{1,1} = p_{0,1} = 0$.

4 Structure of Γ

In this section we consider the structure of Γ and show that if there are multiple sets in the maximal uncoupled partition of Γ, then the measure induced by this Γ cannot be the invariant measure. The next theorem states the main result of this section.

Theorem 2. *Consider the random walk P and its invariant measure m. If m is induced by $\Gamma \subset C$, then Γ is pairwise-coupled.*

The proof of the theorem is deferred to the end of this section. We first introduce some additional notation. For any set Γ_i from the maximal horizontally uncoupled partition of Γ, let

$$B^h(\Gamma_i) = \sum_{(\rho,\sigma)\in\Gamma_i} \alpha(\rho,\sigma)[\sum_{s=-1}^{1} \left(\rho^{1-s}h_s + \rho^{1-s}\sigma p_{s,-1}\right) - \rho], \tag{9}$$

For any set Γ_j from the maximal vertically uncoupled partition of Γ, let

$$B^v(\Gamma_j) = \sum_{(\rho,\sigma)\in\Gamma_j} \alpha(\rho,\sigma)[\sum_{t=-1}^{1} \left(\sigma^{1-t}v_t + \rho\sigma^{1-t}p_{-1,t}\right) - \sigma]. \tag{10}$$

Note that $\sum_{i=1}^{H}\sum_{(\rho,\sigma)\in\Gamma_i} \rho^{i-1}B^h(\Gamma) = 0$ and $\sum_{j=1}^{V}\sum_{(\rho,\sigma)\in\Gamma_j} \sigma^{j-1}B^v(\Gamma) = 0$ are the balance equations for the measure induced by Γ at the horizontal and vertical boundary respectively.

Lemma 2. *Consider the random walk P and a finite measure m induced by some $\Gamma \subset C$. The sequences $\{B^h(\Gamma_i^h)\}_{i=1}^H$ and $\{B^v(\Gamma_i^v)\}_{i=1}^V$ are absolutely convergent.*

Proof. We only provide the proof that the sequence $\{B^h(\Gamma_i^h)\}_{i=1}^H$ is absolutely convergent. We have

$$\begin{aligned}
\sum_{i=1}^{H} |B^h(\Gamma_i^h)| &= \sum_{i=1}^{H} \left| \sum_{(\rho,\sigma)\in\Gamma_i^h} \alpha(\rho,\sigma)[\sum_{s=-1}^{1} \left(\rho^{1-s}h_s + \rho^{1-s}\sigma p_{s,-1}\right) - \rho \right| \\
&\leq \sum_{i=1}^{H} \sum_{(\rho,\sigma)\in\Gamma_i^h} |\alpha(\rho,\sigma)|[\sum_{s=-1}^{1} \left(\rho^{1-s}h_s + \rho^{1-s}\sigma p_{s,-1}\right) + \rho] \\
&< M \sum_{(\rho,\sigma)\in\Gamma} |\alpha(\rho,\sigma)| \frac{1}{1-\rho}\frac{1}{1-\sigma} \\
&< \infty,
\end{aligned}$$

for some $M < \infty$, where the last inequality follows from the fact that m is a finite measure. The result for $\{B^v(\Gamma_i^v)\}_{i=1}^V$ follows similarly. ☐

Remark 1. *Note that Lemma 2 does not require m to be the invariant measure of P, it can be any finite measure.*

The following lemma is a key element for the proof of Theorem 2.

Lemma 3. *Consider the random walk P and a measure m induced by $\Gamma \subset C$. Then m is the invariant measure of P if and only if for all $1 \le i \le H$, $1 \le j \le V$, $B^h(\Gamma^h_i) = 0$ and $B^v(\Gamma^v_j) = 0$.*

Proof. Since m is the invariant measure of P, it satisfies the balance equations at state $(i, 0)$. Therefore,

$$\begin{aligned} 0 &= \sum_{k=-1}^{1} \big[m(i-k,0)h_k + m(i-k,1)p_{k,-1}\big] - m(i,0) \\ &= \sum_{(\rho,\sigma)\in\Gamma} \alpha(\rho,\sigma)\Big[\sum_{k=-1}^{1} \big(\rho^{i-k}h_k + \rho^{i-k}\sigma p_{k,-1}\big) - \rho^i\Big] \qquad (11) \\ &= \sum_{s=1}^{H} \varrho(\Gamma^h_s)^{i-1} \sum_{(\rho,\sigma)\in\Gamma^h_s} \alpha(\rho,\sigma)\Big[\sum_{k=-1}^{1} \big(\rho^{1-k}h_k + \rho^{1-k}\sigma p_{k,-1}\big) - \rho\Big] \\ &= \sum_{s=1}^{H} \varrho(\Gamma^h_s)^{i-1} B^h(\Gamma^h_s). \qquad (12) \end{aligned}$$

The exchange of summations is justified by Lemma 2.

Suppose that H is finite. From (12) it follows that $B^h(\Gamma^h_i)$, $1 \le i \le H$, satisfy a Vandermonde type system of equations. Moreover, from the properties of a maximal horizontally uncoupled partition, the coefficients $\varrho(\Gamma^h_i)$ are all distinct. It follows that $B^h(\Gamma^h_i) = 0$, $1 \le i \le H$. For countably infinite H we resort to [4, Theorem 1], which can be applied based on Lemma 2. Using the same reasoning it follows that $B^v(\Gamma^v_i) = 0$, $1 \le i \le V$, finishing one direction of the proof.

Another direction of the proof can be readily verified by observing that, if $B^h(\Gamma^h_i) = 0$, then $\sum_{i=1}^{H} \rho_i^s B^h(\Gamma^h_i) = 0$ where $s = 0, 1, 2 \cdots$. Therefore, the balance equation for $(i, 0)$, $i > 0$ is satisfied. Using the same reasoning balance at the vertical states is satisfied. Balance in the interior is satisfied by the assumption that m is induced by $\Gamma \subset C$. Finally, balance in the origin is implied by the balance in other parts of the state space. □

Next we present the proof of Theorem 2.

Proof of Theorem 2. The sets of the maximal uncoupled partition can be obtained by taking the union of elements from $\{\Gamma^h_i\}_{i=1}^{H}$ or $\{\Gamma^v_j\}_{j=1}^{V}$. For any

Γ^u_k where $k = 1, \ldots, U$, we can find $I_k \subset \{1, \ldots, H\}$ and $J_k \subset \{1, \ldots, V\}$ such that $\Gamma^u_k = \bigcup_{i \in I_k} \Gamma^h_i = \bigcup_{j \in J_k} \Gamma^v_j$. Using the maximal uncoupled partition, we can introduce measures m_k, defined as

$$m_k(i,j) = \sum_{(\rho,\sigma)\in\Gamma^u_k} \alpha(\rho,\sigma)\rho^i\sigma^j. \tag{13}$$

This allows us to write $m(i,j) = \sum_{k=1}^{U} m_k(i,j)$.[2]

We will show that if measure m is an invariant measure of the random walk in the quarter-plane, then the measures m_k, $k = 1, \ldots, U$, will satisfy all balance equations. Let measure m_k be induced by Γ_k. By definition of C, this implies that all m_k, $k = 1, \ldots, U$, satisfy the balance equations for the states in the interior. Consider the balance equation for m_k at state $(i, 0)$, we will obtain

$$\begin{aligned}
&\sum_{s=-1}^{1} [m_k(i-s,0)h_s + m_k(i-s,1)p_{s,-1}] - m_k(i,0) \\
&= \sum_{s=-1}^{1} \Big[\sum_{(\rho,\sigma)\in\Gamma^u_k} \alpha(\rho,\sigma)\rho^{i-s}h_s + \sum_{(\rho,\sigma)\in\Gamma^u_k} \alpha(\rho,\sigma)\rho^{i-s}\sigma p_{s,-1}\Big] - \sum_{(\rho,\sigma)\in\Gamma^u_k} \alpha(\rho,\sigma)\rho^i \\
&= \sum_{(\rho,\sigma)\in\Gamma^u_k} \alpha(\rho,\sigma)\Big[\sum_{s=-1}^{1} \left(\rho^{i-s}h_s + \rho^{i-s}\sigma p_{s,-1}\right) - \rho^i\Big] \\
&= \sum_{l\in I_k} \varrho(\Gamma^h_l)^{i-1} \sum_{(\rho,\sigma)\in\Gamma^h_l} \alpha(\rho,\sigma)\Big[\sum_{s=-1}^{1} \left(\rho^{1-s}h_s + \rho^{1-s}\sigma p_{s,-1}\right) - \rho\Big] \\
&= \sum_{l\in I_k} \varrho(\Gamma^h_l)^{i-1} B^h(\Gamma^h_l) \\
&= 0.
\end{aligned}$$

By Lemma 2, the interchange of the summations leading to the second equality is valid. The last equality follows from the assumption that m is an invariant measure and Lemma 3.

---

[2] Since the coefficients $\alpha(\rho,\sigma)$ are not necessarily positive it is possible that $m_k(i,j)$ is negative. It is readily verified that in that case m_k is the (signed) measure that is negative on the whole state space. We will see shortly that in that case we can use $-m_k$. Hence we do not introduce the notion of a signed measure.

In similar fashion it follows that the vertical balance equations of m_k are satisfied as well. As a consequence, we have shown that $m_1, \cdots, m_U$ are invariant measures of P. Therefore, if $U > 1$ we have a contradiction to the fact that the invariant measure of an ergodic Markov process is unique up to a constant. □

5 Cardinality of Γ and signs of coefficients

From the last section we know that if the invariant measure m of the random walk P is induced by Γ, then the maximal uncoupled partition of Γ contains only one set and is hence pairwise-coupled. In this section we present conditions on the cardinality of Γ and the signs of the coefficients in the measure induced by Γ.

Theorem 3. *Consider the random walk P and its invariant measure m, where $m(i,j) = \sum_{(\rho,\sigma)\in\Gamma} \alpha(\rho,\sigma)\rho^i\sigma^j$, $\Gamma \subset C$, $\alpha(\rho,\sigma) \neq 0$. Then i) $|\Gamma| \neq 2$ and ii) if $|\Gamma| < \infty$, at least one $\alpha(\rho,\sigma)$ is negative .*

The proof of Theorem 3 follows as a consequence of two lemmas that are given in the remainder of this section. Lemma 4 will prove that $|\Gamma| \neq 2$. Lemma 7 demonstrates necessity of negative coefficients for $|\Gamma| < \infty$.

Lemma 4. *Consider the random walk P and its invariant measure m. If m is induced by a pairwise-coupled set $\Gamma \subset C$, then $|\Gamma| \neq 2$.*

Proof. Without loss of generality, let

$$m(i,j) = \alpha(\rho,\sigma)\rho^i\sigma^j + \alpha(\rho,\tilde{\sigma})\rho^i\tilde{\sigma}^j, \tag{14}$$

where $(\rho,\sigma) \in C$ and $(\rho,\tilde{\sigma}) \in C$.

It follows from the definition of C that σ and $\tilde{\sigma}$ are the roots of the following quadratic equation in x,

$$\sum_{k=-1}^{1}\sum_{s=-1}^{1} \rho^{-s} p_{s,k} x^{1-k} - x = 0. \tag{15}$$

Note that the maximal vertically uncoupled partition of $\{(\rho,\sigma),(\rho,\tilde{\sigma})\}$ consists of the two singleton components $\{(\rho,\sigma)\}$ and $\{(\rho,\tilde{\sigma})\}$. It follows from Lemma 3 that $B^v(\{(\rho,\sigma)\}) = B^v(\{(\rho,\tilde{\sigma})\}) = 0$. Therefore, σ and $\tilde{\sigma}$ are the roots of the following quadratic equation as well

$$\sum_{s=-1}^{1} (\rho p_{-1,s} + v_s)(x^{1-s} - x) = 0. \tag{16}$$

From a comparison of the coefficients of (15) and (16) it follows that either *a)* one of the roots will be 1, contradicting the definition of set C which is restricted within the unit square, or *b)* the transition probabilities of P are such that P is not irreducible and hence not ergodic. Hence, m cannot be the invariant measure of P. $\square$

Before proving the next part of Theorem 3 we will provide a constructive example of a random walk with a measure induced by Γ, with $|\Gamma| = 3$. Note that for $|\Gamma| = 1$ there are many examples in the literature, see, for instance, [3]. Also, for $|\Gamma| = \infty$ constructive examples exist [1].

Example 1. *Consider the random walk with $p_{-1,1} = 2/5$, $p_{0,-1} = 2/5$, $p_{1,-1} = 1/5$, $h_1 = 1/5$, $h_0 = 4/5$, $v_{-1} = 18/25$, $v_0 = 7/25$ and all other probabilities zero. Also, consider the measure $m(i,j) = \sum_{k=1}^{3} \alpha_k \rho_k^i \sigma_k^j$, where $(\rho_1, \sigma_1) = (1/2, 1/4)$, $(\rho_2, \sigma_2) = (1/16, 1/4)$, $(\rho_3, \sigma_3) = (1/16, 1/36)$, $\alpha_1 = 1$, $\alpha_2 = -20/7$ and $\alpha_3 = 862/231$. It can be readily verified that $m(i,j)$ satisfies all balance equations, hence that $m(i,j)$ is the invariant measure of the random walk.*

We now proceed to proving the next part of Theorem 3. We introduce a final piece of notation and two technical results that will help in the proofs. Let

$$b^h(\Gamma_i^h) = \frac{B^h(\Gamma_i^h)}{\varrho(\Gamma_i^h) \sum_{(\rho,\sigma)\in\Gamma_i} \alpha(\rho,\sigma)} + \left(1 - \frac{1}{\varrho(\Gamma_i^h)}\right) h_1 + \left(1 - \varrho(\Gamma_i^h)\right) h_{-1} \tag{17}$$

and

$$b^v(\Gamma_j^v) = \frac{B^v(\Gamma_j^v)}{\varsigma(\Gamma_j^v) \sum_{(\rho,\sigma)\in\Gamma_j} \alpha(\rho,\sigma)} + \left(1 - \frac{1}{\varsigma(\Gamma_j^v)}\right) v_1 + \left(1 - \varsigma(\Gamma_j^v)\right) v_{-1}. \tag{18}$$

Lemma 5. *If $\tilde{\sigma} > \sigma$, $\tilde{\rho} > \rho$ and $\alpha(\rho,\sigma) > 0$ then*

$$b^h(\{(\rho,\sigma),(\rho,\tilde{\sigma})\}) > b^h(\{(\rho,\sigma)\}), \quad b^h(\{(\rho,\sigma),(\rho,\tilde{\sigma})\}) < b^h(\{(\rho,\tilde{\sigma})\}),$$
$$b^v(\{(\rho,\sigma),(\tilde{\rho},\sigma)\}) > b^v(\{(\rho,\sigma)\}), \quad b^v(\{(\rho,\sigma),(\tilde{\rho},\sigma)\}) < b^v(\{(\tilde{\rho},\sigma)\}).$$

Proof. From the definition in (17) it follows that

$$\begin{aligned} b^h(\{(\rho,\sigma),(\rho,\tilde{\sigma})\}) =& \frac{\alpha(\rho,\sigma)\sigma+\alpha(\rho,\tilde{\sigma})\tilde{\sigma}}{\alpha(\rho,\sigma)+\alpha(\rho,\tilde{\sigma})}(\rho p_{-1,-1}+p_{0,-1}+\frac{1}{\rho}p_{1,-1}) - \\ & p_{1,1}-p_{0,1}-p_{-1,1}, \end{aligned}$$

$$b^h(\{(\rho,\sigma)\}) = \sigma(\rho p_{-1,-1}+p_{0,-1}+\frac{1}{\rho}p_{1,-1})-p_{1,1}-p_{0,1}-p_{-1,1},$$

and

$$b^h(\{(\rho,\tilde{\sigma})\}) = \tilde{\sigma}(\rho p_{-1,-1}+p_{0,-1}+\frac{1}{\rho}p_{1,-1})-p_{1,1}-p_{0,1}-p_{-1,1}.$$

From the above the first row of inequalities follow directly. The remaining inequalities follow directly from (18). ☐

The second technical lemma that we will need is readily verified and stated here without proof.

Lemma 6. *If* $t_1(1-\rho)+t_2(1-\tilde{\rho}) \geq 0$, $t_1(1-1/\rho)+t_2(1-1/\tilde{\rho}) \geq 0$ *and* $\tilde{\rho} > \rho$, *then* $t_1 \leq 0$ *and* $t_2 \geq 0$.

The second part of Theorem 3 for $|\Gamma| < \infty$ follows from the next lemma.

Lemma 7. *Consider the random walk* P *and its invariant measure* m, *where* $m(i,j) = \sum_{(\rho,\sigma)\in\Gamma} \alpha(\rho,\sigma)\rho^i\sigma^j$, $\Gamma \subset C$, $\alpha(\rho,\sigma) \neq 0$. *If* $|\Gamma| < \infty$, *then at least one* $\alpha(\rho,\sigma)$ *is negative.*

Proof. Without loss of generality we assume that $\rho_1 = \rho_2$, $\sigma_1 > \sigma_2$ and $\sigma_2 \geq \sigma_3 \geq \cdots \geq \sigma_n$. Moreover, $\varsigma(\Gamma_j^V)$ is strictly decreasing in j in the following two cases which will be analyzed separately. The first case is $\rho_2 > \rho_3$, $\rho_3 \geq \rho_4 \geq \cdots \geq \rho_n$ and $\varrho(\Gamma_i^H)$ is strictly decreasing in i. The second case is $\rho_2 < \rho_3$, $\rho_3 \leq \rho_4 \leq \cdots \leq \rho_n$ and $\varrho(\Gamma_i^H)$ is strictly increasing with i.

For the first case we consider the relations

$$\begin{aligned} (1-1/\rho_1)\,h_1+(1-\rho_1)h_{-1} &= b^h(\Gamma_1^h), \\ (1-1/\rho_n)\,h_1+(1-\rho_n)h_{-1} &= b^h(\Gamma_H^h), \\ (1-1/\sigma_1)\,v_1+(1-\sigma_1)v_{-1} &= b^v(\Gamma_1^v), \\ (1-1/\sigma_n)\,v_1+(1-\sigma_n)v_{-1} &= b^v(\Gamma_V^v), \end{aligned} \tag{19}$$

which by Lemma 3 are required to hold if m is the invariant measure of the random walk P. We will construct s_1, s_2, t_1 and t_2 that satisfy

$$\begin{aligned} (1-1/\rho_1)\, s_1 + (1-1/\rho_n)\, s_2 &\geq 0, \\ (1-\rho_1)\, s_1 + (1-\rho_n)\, s_2 &\geq 0, \\ (1-1/\sigma_1)\, t_1 + (1-1/\sigma_n)\, t_2 &\geq 0, \\ (1-\sigma_1)\, t_1 + (1-\sigma_n)\, t_2 &\geq 0 \end{aligned} \tag{20}$$

and

$$b^h(\Gamma^h_1)s_1 + b^h(\Gamma^h_H)s_2 + b^v(\Gamma^v_1)t_1 + b^v(\Gamma^v_V)t_2 < 0. \tag{21}$$

By Farkas' Lemma this leads to a contradiction to (19) because the transition probabilities h_1, h_{-1}, v_1, v_{-1} are non-negative. The s_1, s_2, t_1 and t_2 are constructed by considering the auxiliary measure $\bar{m} = \alpha(\rho_1, \sigma_1)\rho_1^i\sigma_1^j + \alpha(\rho_n, \sigma_n)\rho_n^i\sigma_n^j$ and the two-dimensional random walk $\bar{P}$, that has the same transition probabilities as P in the interior of the state space and transition probabilities $\bar{h}_1$, $\bar{h}_{-1}$, $\bar{v}_1$ and $\bar{v}_{-1}$ along the boundaries. We now consider the relations

$$\begin{aligned} (1-1/\rho_1)\, \bar{h}_1 + (1-\rho_1)\bar{h}_{-1} &= b^h(\{(\rho_1, \sigma_1)\}), \\ (1-1/\rho_n)\, \bar{h}_1 + (1-\rho_n)\bar{h}_{-1} &= b^h(\{(\rho_n, \sigma_n)\}), \\ (1-1/\sigma_1)\, \bar{v}_1 + (1-\sigma_1)\bar{v}_{-1} &= b^v(\{(\rho_1, \sigma_1)\}), \\ (1-1/\sigma_n)\, \bar{v}_1 + (1-\sigma_n)\bar{v}_{-1} &= b^v(\{(\rho_n, \sigma_n)\}). \end{aligned} \tag{22}$$

If (22) would hold for $\bar{h}_1 = h_1$, $\bar{h}_{-1} = h_{-1}$, $\bar{v}_1 = v_1$ and $\bar{v}_{-1} = v_{-1}$, $\bar{m}$ would be the invariant measure of P which contradicts the assumption that m is the invariant measure of P. However, if $\bar{P}$ is ergodic (22) cannot hold due to Theorem 2. If $\bar{P}$ is not ergodic, there is no finite invariant measure and (22) cannot hold either. Therefore, (22) is not satisfied for any non-negative $\bar{h}_1$, $\bar{h}_{-1}$, $\bar{v}_1$ and $\bar{v}_{-1}$. By Farkas' Lemma, there exist s_1, s_2, t_1 and t_2 that satisfy (20) and

$$b^h(\{(\rho_1, \sigma_1)\})s_1 + b^h(\{(\rho_n, \sigma_n)\})s_2 + b^v(\{(\rho_1, \sigma_1)\})t_1 + b^v(\{(\rho_n, \sigma_n)\})t_2 < 0.$$

Note, that from Lemma 5 it follows that $b^h(\{\Gamma^h_1\}) < b^h(\{(\rho_1, \sigma_1)\})$,

$b^h(\{\Gamma^h_H\}) \geq b^h(\{(\rho_n, \sigma_n)\})$, $b^v(\{\Gamma^v_1\}) = b^v(\{(\rho_1, \sigma_1)\})$ and $b^v(\{\Gamma^v_V\}) \geq b^v(\{(\rho_n, \sigma_n)\})$. Also, from Lemma 6 it follows that $s_1 \geq 0$, $s_2 \leq 0$, $t_1 \geq 0$, $t_2 \leq 0$. Therefore, s_1, s_2, t_1 and t_2 satisfy (21). This concludes the proof of the first case.

For the second case we consider the relations

$$\begin{aligned} (1-1/\sigma_1)\, v_1 + (1-\sigma_1) v_{-1} &= b^v(\Gamma_1^v), \\ (1-1/\sigma_2)\, v_1 + (1-\sigma_2) v_{-1} &= b^v(\Gamma_2^v), \end{aligned} \tag{23}$$

that are necessary for m to be the invariant measure and obtain a contradiction by constructing t_1 and t_2 that satisfy

$$(1-1/\sigma_1)\, t_1 + (1-1/\sigma_2)\, t_2 \geq 0, \tag{24}$$

$$(1-\sigma_1)\, t_1 + (1-\sigma_2)\, t_2 \geq 0, \tag{25}$$

$$b^v(\Gamma_1^v) t_1 + b^v(\Gamma_1^v) t_2 < 0. \tag{26}$$

The auxiliary measure that is used is $\tilde{m}(i,j) = \alpha(\rho_1,\sigma_1)\rho_1^i\sigma_1^j + \alpha(\rho_2,\sigma_2)\rho_2^i\sigma_2^j$. Observe that $\rho_1 = \rho_2$ and that the corresponding relations are

$$\begin{aligned} (1-1/\rho_1)\, h_1 + (1-\rho_1) h_{-1} &= b^h(\{(\rho_1,\sigma_1),(\rho_2,\sigma_2)\}), \\ (1-1/\sigma_1)\, v_1 + (1-\sigma_1) v_{-1} &= b^v(\{(\rho_1,\sigma_1)\}), \\ (1-1/\sigma_2)\, v_1 + (1-\sigma_2) v_{-1} &= b^v(\{(\rho_2,\sigma_2)\}). \end{aligned}$$

From Farkas' Lemma and Lemma 4 it follows that there exist s_1, t_1 and t_2 that satisfy (24), (25) and

$$b^h(\{(\rho_1,\sigma_1),(\rho_2,\sigma_2)\}) s_1 + b^v(\{(\rho_1,\sigma_1)\}) t_1 + b^v(\{(\rho_2,\sigma_2)\}) t_2 \leq 0, \tag{27}$$

where $s_1 = 0$, since it satisfies $(1-1/\rho_1)s_1 \geq 0$ and $(1-\rho_1)s_1 \geq 0$. Moreover, since, $b^v(\Gamma_1^v) = b^v(\{(\rho_1,\sigma_1)\})$ and, by Lemma 5, we have $b^v(\Gamma_2^v) > b^v(\{(\rho_2,\sigma_2)\})$. Moreover, by Lemma 6 we have, $t_1 \geq 0, t_2 \leq 0$. Then it follows that t_1 and t_2 satisfy (26). This concludes the proof of the second case. □

6 Acknowledgments

The authors wish to thank Anton Stoorvogel for some useful discussions. Moreover they would like to thank Bert Zwart and the associate editor for some pointers to existing literature. Yanting Chen gratefully acknowledges support through a CSC scholarship[No.2008613008]. This work was supported in part by NWO grant 612.001.107.

References

[1] I. Adan, G.J. van Houtum, J. Wessels, and W.H.M. Zijm. A compensation procedure for multiprogramming queues. *OR Spectrum*, 15(2):95–106, 1993.

[2] I. Adan, J. Wessels, and W.H.M Zijm. A compensation approach for two-dimensional Markov processes. *Advances in Applied Probability*, 25(4):783–817, 1993.

[3] R.J. Boucherie and N.M. van Dijk, editors. *Queueing Networks: A Fundamental Approach.* Springer Verlag, 2010.

[4] L. Brown, A. Shields, and K. Zeller. On absolutely convergent exponential sums. *Transactions of the American Mathematical Society*, 96(1):162–183, 1960.

[5] J.W. Cohen and O.J. Boxma. *Boundary value problems in queueing system analysis.* North Holland, 1983.

[6] A.B. Dieker and J. Moriarty. Reflected Brownian motion in a wedge: sum-of-exponential stationary densities. *Electron. Commun. Probab*, 14:1–16, 2009.

[7] G. Fayolle, R. Iasnogorodski, and V.A. Malyshev. *Random walks in the quarter-plane: algebraic methods, boundary value problems and applications.* Springer Verlag, 1999.

[8] M. Miyazawa. Light tail asymptotics in multidimensional reflecting processes for queueing networks. *Top*, 1–67, 2011.

[9] M.F. Neuts. *Matrix-geometric solutions in stochastic models: an algorithmic approach.* Dover Publications, 1981.